

Facilitated diffusion of DNA-binding proteins: Simulation of large systems

Holger Merlitz,^{1,*} Konstantin V. Klenin,² Chen-Xu Wu,^{1,†} and Jörg Langowski²

¹*Softmatter Lab, Department of Physics, Xiamen University, Xiamen 361005, P.R. China*

²*Division of Biophysics of Macromolecules, German Cancer Research Center, D-69120 Heidelberg, Germany*

(Dated: February 2, 2008)

The recently introduced method of excess collisions (MEC) is modified to estimate diffusion-controlled reaction times inside systems of arbitrary size. The resulting MEC-E equations contain a set of empirical parameters, which have to be calibrated in numerical simulations inside a test system of moderate size. Once this is done, reaction times of systems of arbitrary dimensions are derived by extrapolation, with an accuracy of 10 to 15 percent. The achieved speed up, when compared to explicit simulations of the reaction process, is increasing proportional to the extrapolated volume of the cell.

PACS numbers: 87.16.Ac

1. INTRODUCTION

Diffusion controlled bio-chemical reactions play a central role in keeping any organism alive [1, 2]: The transport of molecules through cell membranes, the passage of ions across the synaptic gap, or the search carried out by drugs on the way to their protein receptors are predominantly diffusive processes. Further more, essentially all of the biological functions of DNA are performed by proteins that interact with specific DNA sequences [3, 4], and these reactions are diffusion-controlled.

However, it has been realized that some proteins are able to find their specific binding sites on DNA much more rapidly than is ‘allowed’ by the diffusion limit [1, 5]. It is therefore generally accepted that some kind of facilitated diffusion must take place in these cases. Several mechanisms, differing in details, have been proposed. All of them essentially involve two steps: the binding to a random non-specific DNA site and the diffusion (sliding) along the DNA chain. These two steps may be reiterated many times before proteins actually find their target, since the sliding is occasionally interrupted by dissociation. Berg [5] and Zhou [6] have provided thorough (but somewhat sophisticated) theories that allow estimates for the resulting reaction rates. Recently, Halford and Marko have presented a comprehensive review on this subject and proposed a remarkably simple and semi-quantitative approach that explicitly contains the mean sliding length as a parameter of the theory [7]. This approach has been refined and put onto a rigorous base in a recent work by the authors [8]. A plethora of scaling regimes have been studied for a large range of chain densities and protein-chain affinities in a recent work by Hu et al. [9].

The numerical treatment of such a reaction is efficiently done with the method of excess collisions [10] (MEC), where the reverse process (protein departs from

the binding site and propagates toward the periphery of the cell) is simulated. This approach delivers exact results and a significant speed up when compared to straight forward simulations. Unfortunately, once very large systems are under investigation, the numerical treatment of the DNA chain (whose length is proportional to the volume of the cell) quickly turns into a bottleneck, since the MEC approach requires the construction of the cell in its full extent. Realistic cell models have to deal with thermal fluctuations of the chain and its hydrodynamic interaction, thereby imposing a strict limit to the size that can be managed. In the present work we demonstrate how to implement a modification of the MEC approach that allows to simulate a test system of reasonable size, followed by an extrapolation to cells of arbitrary size.

After a definition of the problem in Sect. 2.1, the MEC approach is briefly summarized in Sect. 2.2. In Sect. 2.3 the numerical implementation of facilitated diffusion is presented, and 2.4 delivers an analytical estimate for the reaction time. As a preparation for the random walk simulations, the chain is constructed in Sect. 3 and the specific recurrence times are evaluated inside a small test system (Sect. 4). In Sect. 5, random walk simulations are carried out in order to construct the empirical MEC-E equations. These are then employed to extrapolate the reaction times to cells of much larger dimensions in Sect. 6. A comparison with exact solutions (in the case of free diffusion) and the analytical estimate of Sect. 2.4 suggests that the MEC-E approach delivers an accuracy of 10 to 15 percent with a speed up of several orders of magnitude.

2. METHODOLOGY

2.1. Definition of the system

As a *cell* we define a spherical volume of radius R , containing a chain (‘DNA’) of length L and a specific binding target of radius R_a . The target is located in the middle of the chain, that in turn coincides with the cen-

*Electronic address: merlitz@gmx.de

†Electronic address: cxwu@jingxian.xmu.edu.cn

ter of the cell. The state of the system is well defined with the position of a random walker ('protein'), which can either diffuse freely inside the cell or, temporarily, associate with the chain to propagate along the chain's contour (the numerical realization of this process is discussed in detail in Sect. 2.3). The distance of the walker from the center defines the (radial) reaction coordinate r . We shall further denote the periphery of the central target (at $r = R_a$) as **state A** and the periphery of the cell ($r = R$) as **state B**. To be investigated is the average reaction time τ_{BA} the walker needs to propagate from **B** to **A** as a function of the binding affinity between walker and chain.

2.2. Method of excess collisions (MEC)

The MEC approach was presented in its full generality elsewhere [10, 11]. In short, it allows to determine the reaction time τ_{BA} while simulating the back reaction **A** $\rightarrow$ **B** (average reaction time: τ_{AB}) using the relation

$$\tau_{BA} = (N_{\text{coll}} + 1) \cdot \tau_R - \tau_{AB} . \quad (1)$$

The walker starts at the center ($r(t=0) = 0$) and propagates towards the periphery ($r(t = \tau_{AB}) = R$), a process that is much faster than its reversal ($\tau_{AB} \ll \tau_{BA}$). On its way to **B**, the walker may repeatedly return back to **A**; such an event is called collision, and N_{coll} stands for the average number of collisions. τ_R is the recurrence time and evaluated via

$$\tau_R = \tilde{\tau}_R V_{\text{eff}}(R) , \quad (2)$$

where we have defined the specific recurrence time

$$\tilde{\tau}_R \equiv \frac{\tau_R^*}{V_{\text{eff}}(R_a)} , \quad (3)$$

a quantity, which is derived from simulations of the recurrence time τ_R^* within a small test system of the size of the central target (Sect. 4). The effective volume is defined as

$$V_{\text{eff}} \equiv \int_V \exp \left[\frac{-U(\mathbf{r})}{k_B T} \right] d\mathbf{r} , \quad (4)$$

and depends upon the energy of the walker $U(\mathbf{r})$ and hence the implementation of the binding potential between walker and chain.

2.3. Simple model for facilitated diffusion of DNA-binding proteins

The nonspecific binding of the walker to the chain is accounted for by the attractive step potential

$$U(s) = \begin{cases} -E_o & s \leq r_c \\ 0 & s > r_c , \end{cases} \quad (5)$$

where s is the shortest distance between walker and chain. This defines a pipe with radius r_c around the chain contour that the walker is allowed to enter freely from outside, but to exit only with the probability

$$p = \exp(-E_o/k_B T) , \quad (6)$$

where $k_B T$ is the Boltzmann factor, otherwise it is reflected back inside the chain. We may therefore denote p as *exit probability*. This quantity allows to define the equilibrium constant K of the two phases, the free and the non-specifically bound protein, according to

$$K \equiv \frac{\sigma}{c} = \frac{V_c}{L} \left(\frac{1}{p} - 1 \right) , \quad (7)$$

where c is the concentration of free proteins and σ the linear density of non-specifically bound proteins. $V_c = \pi r_c^2 L$ is the geometric volume of the chain. It should be noted that in our previous publication [10], σ was defined as $\sigma = c V_c / (p L)$, with the disadvantage of being non-zero in case of vanishing protein-chain interaction ($p = 1$). The present choice defines σ as the excess concentration of proteins along the chain contour and leads to a vanishing sliding-length (Eq. 14) in case of free diffusion.

The specific binding site is a spherical volume, located in the middle of the chain and of identical radius, i.e. $R_a = r_c$. Applying the walker-chain potential Eq. (5), the effective volume Eq. (4) of the cell becomes

$$V_{\text{eff}}(R) = V + V_c \left(\frac{1}{p} - 1 \right) , \quad (8)$$

and that of the central target is simply

$$V_{\text{eff}}(R_a) = \frac{V_a}{p} = \frac{4\pi R_a^3}{3p} . \quad (9)$$

2.4. Analytical estimate for the reaction time and definition of the sliding length

In case of free diffusion and for a spherical cell, Szabo et al. have evaluated the exact solution for the time a walker needs to reach the radius R_a , after starting at the periphery R , yielding [12]

$$\tau_{Sz} = \frac{R^2}{3D} \cdot \left(\frac{R}{R_a} + \frac{R_a^2}{2R^2} - \frac{3}{2} \right) . \quad (10)$$

Here, D is the diffusion coefficient. In presence of the chain, exact solutions are known for simple geometrical setups only [5], but as discussed elsewhere [8], it is still possible to approximate the reaction time using an analytical approach, once certain conditions are satisfied. The resulting expression is

$$\tau_{BA}(\xi) = \left(\frac{V}{8D_{3d}\xi} + \frac{\pi L \xi}{4D_{1d}} \right) \left[1 - \frac{2}{\pi} \arctan \left(\frac{R_a}{\xi} \right) \right] \quad (11)$$

with the 'sliding' variable

$$\xi = \sqrt{\frac{D_{1d} K}{2\pi D_{3d}}} \quad (12)$$

and D_{1d} and D_{3d} being the diffusion coefficients in sliding-mode and free diffusion, respectively. Generally, the equilibrium constant K has to be determined in simulations of a (small) test system, containing a piece of chain without specific binding site. In the present model, K is known analytically via Eq. (7). If the step-size dr of the random walker is equal both inside and outside the chain (the direction of the step being arbitrary), we further have

$$D_{1d} = D_{3d} = \frac{dr^2}{6}, \quad (13)$$

and hence obtain

$$\xi = \sqrt{\frac{r_c^2}{2} \left(\frac{1}{p} - 1 \right)}. \quad (14)$$

This variable has got the dimension of length; as we have pointed out in [8], it corresponds to the average *sliding length* of the protein along the DNA contour in the model of Halford and Marko [7] and we shall henceforth use the same expression for ξ . In case of free diffusion ($p = 1$), the sliding length is zero and Eq. (11) simplifies to

$$\tau_{BA}(\xi = 0) = \frac{R^3}{3 R_a D_{3d}}, \quad (15)$$

which equals Szabo's result Eq. (10) in leading order of R/R_a .

3. NUMERICAL MODEL

In order to approximate the real biological situation, the DNA was modeled as a chain of straight segments of equal length l_0 . Its mechanical stiffness was introduced in terms of a bending energy associated with each chain joint:

$$E_b = k_B T \alpha \theta^2, \quad (16)$$

where α represents the dimensionless stiffness parameter, and θ the bending angle. The numerical value of α defines the persistence length (l_p), i.e. the "stiffness" of the chain. The excluded volume effect was taken into account by introducing the effective chain radius r_c . The conformations of the chain, with distances between non-adjacent segments smaller than $2r_c$, were forbidden. The target of specific binding was assumed to lie exactly in the middle of the DNA. The whole chain was packed in a spherical volume (cell) of radius R in such a way that the target occupied the central position.

To achieve a close packing of the chain inside the cell, we used the following algorithm. First, a relaxed conformation of the free chain was produced by the standard




FIG. 1: Upper part: 2-dimensional projection of a 3-dimensional random chain-contour of length $L = 400.2$ (persistence lengths) confined inside a spherical cell of radius $R = 6$. Lower part: Radial chain density distribution, averaged over 20 conformations. Beyond $r = 4$ (dashed line), the density declines rapidly.

Metropolis Monte-Carlo (MC) method. For the further compression, we defined the center-norm (c-norm) as the maximum distance from the target (the middle point) to the other parts of the chain. Then, the MC procedure was continued with one modification. Namely, a MC step was rejected if the c-norm was exceeding 105% of the lowest value registered so far. The procedure was stopped when the desired degree of compaction was obtained.

Below in this paper, one step dt was chosen as the unit of time and one persistence length $l_p = 50$ nm of the DNA chain as the unit of distance. The following values of parameters were used. The length of one segment was chosen as $l_0 = 0.2$, so that one persistence length was partitioned into 5 segments. The corresponding value of the stiffness parameter was $\alpha = 2.403$ [13]. The chain radius was $r_c = 0.06$, and the active site was modeled as a sphere of identical radius $r_a = 0.06$ embedded into the chain. The step-size of the random walker both inside and outside the chain was $dr = 0.02$, corresponding to a diffusion coefficient $D_{3d} = D_{1d} = dr^2/6 = 2 \cdot 10^{-4}/3$.

Figure 1 displays a typical chain, and the radial chain density, obtained with Monte Carlo integration and averaged over 20 different chain conformations. The strong increase of chain density towards the center is merely a geometric effect and caused by the chain passing through the origin. Close to the periphery of the cell, the den-

sity was rapidly declining since the contour was forced to bend back inwards. Within a radius of $r < 4$, however, the chain content remained reasonably constant, and the medium could be regarded as approximately homogeneous.

4. COMPUTATION OF THE SPECIFIC RECURRENCE TIME

To compute the specific recurrence time $\tilde{\tau}_R$ of Eq. (3), the recurrence time inside a small test system (here: the central binding target of radius R_a) has to be determined. To achieve that, the entire system, i.e. the spherical target and a short piece of chain, was embedded into a cube of $4R_a$ side-length with reflective walls. In principle, the size of the cube should be of no relevance, but it was found that, if chosen too small, effects of the finite step-size were emerging. The walker started inside the sphere. Each time upon leaving the spherical volume a collision was noted. If the walker was about to exit the cylindrical volume of the chain, it was reflected back inside with the probability $1 - p$ (Eq. 6). The clock was halted as long as the walker moved outside the sphere and only counted time-steps inside the sphere. The resulting recurrence time τ_R^* has to be divided by the effective volume of the central target, Eq. (9), to yield the specific recurrence time $\tilde{\tau}_R$. Table I contains the results for a set of different walker-chain affinities.

5. DIFFUSION INSIDE THE CELL

The goal is to analyze the propagation of the walker within a small cell of radius R_S and to extrapolate the results to a larger system of arbitrary size $R_L > R_S$. As a test site we have set up a cell of radius $R = 6$, containing a chain of length $L = 400.2$ (Figure 1). The walker was starting at the center ($r = 0$) and moving towards the periphery of the cell. Such a process shall be denoted as *run*. Whenever the walker returned back to the binding site ($r < R_a$), one collision was noted. A set of 2000 runs, including 20 different chain conformations, was carried out for each value of the exit parameter p , which is related to the walker-chain affinity via Eq. (6). For a set of reaction coordinates r_i , the first arrival times were monitored, as well as the number of collisions that had occurred before first passage.

5.1. The effective diffusion coefficient

Figure 2 displays the first arrival times (left) for different exit probabilities p . To analyse the diffusive properties of the propagation, the arrival times were fitted using the macroscopic diffusion law

$$\tau_f(p, r) = \frac{r^\alpha}{6D_{\text{eff}}(p)} \quad (17)$$




FIG. 2: First passage times (left) and number of collisions (right) as a function of the reaction coordinate r , for various exit probabilities $p = 2^{-l}$ and $l = 3, 5, 7, 9, 11$ (bottom to top plots). The curves are χ^2 -fits of Eq. (17) (left) and Eq. (18) (right) within the range $\xi < r < 4$ and extrapolated to $r = 6$.

with an effective diffusion coefficient $D_{\text{eff}}(p)$. For low and moderate values of the walker-chain affinity, the arrival times were well described when assuming regular diffusion, i.e. an exponent of $\alpha = 2$. At high walker-chain affinities, this exponent was growing larger, indicating the onset of anomalous subdiffusion. Table I contains the fit parameters when the fits were carried out within the range $\xi < r < 4$, and the solid curves in figure 2 (left) display the resulting functional form of Eq. (17), when extrapolated to the full range $0 < r < 6$.

The lower boundary of the fit range, the sliding length ξ , was implemented because the near the central target, the transport process was dominated by one dimensional sliding rather than three dimensional diffusion. The upper boundary was introduced since the chain distribution beyond $r > 4$ was affected by boundary effects near the periphery of the cell, as is clearly visible at Figure 1. Within the range of $\xi < r < 4$, however, the propagation of the walker could approximately be regarded as a random walk inside a homogeneous and crowded environment.

5.2. The functional dependence of N_{coll} on the target-distance

The right hand side of Figure 2 displays the number of collisions N_{coll} as a function of the radius r for var-

ious walker-chain affinities. Quite generally, there exists a steep increase close to the central target, after which the function gradually levels off to reach a plateau. In Appendix A, we argue that this functional behavior can be described as

$$N_{\text{coll}}(r) = \frac{N_{\infty} \cdot (r - R_{\text{eff}})}{r}, \quad (18)$$

where N_{∞} stands for the asymptotic limit $N_{\text{coll}}(r \rightarrow \infty)$ and R_{eff} defines an effective target size. As a result of facilitated diffusion, the mode of propagation is predominantly one-dimensional near the central target. This relation is therefore invalid within a radius of the average sliding length of the walker and should be applied for $r > \xi$. Under this condition, both N_{∞} and R_{eff} were used as free fit-parameters and the fit range was restricted to $\xi < r < 4$, for the same reason as discussed in Sec. 5.1. The solid curves of Figure 2 (right) display the best fits (extrapolated to $r = 6$), and Table I contains the corresponding values for the fit-parameters.

An alternative approach to $N_{\text{coll}}(r)$ is described in Appendix B, leading to

$$N_{\text{coll}}(r) = \frac{f(r)}{V_{\text{eff}} \tilde{\tau}_R}, \quad (19)$$

where $f(r)$ is defined in Eq. (B2). It contains both parameters D_{eff} and R_{eff} which are used as free fit parameters to determine the effective diffusion coefficient and an effective target size. The results are given in Table I. The effective volume $V_{\text{eff}}(r)$ as a function of radius r is actually a complicated function that depends on the radial chain density (Fig. 1), but for this investigation

we have assumed a perfectly homogeneous chain density and evaluated

$$V_{\text{eff}}(r) = \frac{V(r) V_{\text{eff}}(R)}{V(R)} \quad (20)$$

with the cell-radius $R = 6$. When comparing the best fits for the effective diffusion coefficient with the results of Eq. (17), the agreement is only qualitative. In fact, Eq. (20) does not deliver an accurate way to determine D_{eff} . This may be so because the second term of function

$$f(r) = \frac{r^2}{3 D_{\text{eff}}} \cdot \left(\frac{r}{R_{\text{eff}}} + \frac{R_{\text{eff}}^2}{2 r^2} - 1 \right),$$

the fraction R_{eff}^2/r^2 , quickly drops to zero and hence both fit-parameters D_{eff} and R_{eff} become linear dependent. This implies that D_{eff} is actually determined locally, close to the (effective) target, and not averaged over $\xi < r < 4$. Except for high walker-chain affinities, Eq. (17) delivers a more accurate description of the diffusion process, which is verified with the quadratic dependence of the passage time on the reaction coordinate.

The effective target size R_{eff} agrees fairly well with the corresponding findings of Eq. (18) and increases substantially with the walker-chain affinity. As a consequence of facilitated diffusion, the walker initially moves away from the target in one-dimensional sliding mode, and its (effectively) free diffusion begins further outside, thereby increasing the effective target size. Hence it is no surprise to find R_{eff} being of similar dimension as the average sliding length ξ (Table I).

5.3. The empirical MEC-E equations

It is now possible to combine Equations (18) and (17) with (1) and (2) to obtain the empirical MEC-E equations

$$\tau_{\text{BA}}(p, r) = (N_{\text{coll}}(p, r) + 1) \tilde{\tau}_R V_{\text{eff}} - \tau_f(p, r), \quad (21)$$

which allow to evaluate the reaction time $\tau_{\text{BA}}(p, r)$ for any reaction coordinate r by extrapolation of the number of collisions $N_{\text{coll}}(p, r)$ and the first arrival times $\tau_f(p, r)$.

When using Eq. (19) instead of (18), we obtain

$$\tau_{\text{Sz,eff}}(p, r) = \frac{r^2}{3 D_{\text{eff}}(p)} \cdot \left(\frac{r}{R_{\text{eff}}(p)} + \frac{R_{\text{eff}}^2(p)}{2 r^2} - \frac{3}{2} \right), \quad (22)$$

which can be regarded as an empirical generalization of Szabo's exact result for free diffusion, Eq. (10).

Since both sets of equations are based on the MEC approach, while employing different ways to extrapolate the number of collisions to large cells, we will refer to them as MEC-E equations. In the following section we will apply both approaches, Eq. (21) and Eq. (22), to extrapolate the reaction times to large cell radii, and compare their results.

TABLE I: The first column is the exponent of the exit probability $p = 2^{-l}$, the second column the corresponding sliding parameter, followed by the specific recurrence time (Sect. 4). The next six columns contain optimized parameters of the χ^2 -fits of equations (17), (18) and (19). The last column defines the speed up achieved with the extrapolation from $R_S = 4$ to $R_L = 6$, when compared with the explicit simulation of the reaction time $\tau_{\text{BA}}(R_L)$.

l	ξ	$\tilde{\tau}_R$	$D_{\text{eff}}^{(1)}$	α	R_{eff}	N_{∞}	$D_{\text{eff}}^{(1)}$	R_{eff}	$\frac{\tau_{\text{BA}}(R_L)}{\tau_{\text{BA}}(R_S)}$
Eq.	(14)	(3)	(17)		(18)		(19)		
0	0	4464	6.63	2	0.064	3.83	6.09	0.060	520
1	0.042	2594	6.66	2	0.078	6.42	5.98	0.069	410
2	0.073	1413	6.55	2	0.084	9.95	6.42	0.079	354
3	0.112	741.6	6.35	2	0.100	15.7	6.73	0.092	292
4	0.164	379.7	5.97	2	0.118	25.0	6.42	0.116	221
5	0.236	192.6	5.37	2	0.174	39.7	5.27	0.167	169
6	0.337	96.81	4.50	2	0.224	61.9	4.28	0.231	120
7	0.478	48.62	3.67	2	0.319	95.3	2.95	0.348	94
8	0.677	24.30	2.83	2	0.417	135	2.09	0.491	70
9	0.959	12.17	2.47	2.07	0.59	199	1.28	0.69	61
10	1.357	6.089	2.44	2.20	0.81	279	0.63	1.10	54
11	1.920	3.044	2.41	2.27	1.10	398	0.35	1.50	63

1) In units of 10^{-5}




FIG. 3: Reaction time τ_{BA} of the protein as a function of the sliding length Eq. (14). The explicit simulation (solid dots) required about 140 times the number of simulation steps of the extrapolation using Eq. (21) (open circles) or Eq. (22) (triangles). The curve is the analytical estimate Eq. (11).

6. RESULTS

As a first consistency-check, the MEC-E equations were applied to estimate the reaction time τ_{BA} of the walker entering the cell at radius $R = 6$. The simulation of the reaction $\mathbf{B} \rightarrow \mathbf{A}$ was additionally carried out explicitly and the results are displayed in Figure 3. The effective volume of the cell was evaluated via Eq. (8), using the total chain length $L = 400.2$. The results, shown in Figure 3, imply that the extrapolation from $R_S = 4$ (the radius used to optimize the parameters) to $R_L = 6$ delivered accurate results for the reaction times. This should not be taken for granted, taking into account the problematic chain distribution between $R_S < r < R_L$. In fact, $\tau_f(p, r)$ becomes inaccurate in this region due to anomalous diffusion (Figure 2, left), but this term contributes just a small amount to Eq. (21), since for reasonably large cells the first arrival time τ_f is small compared to the corresponding reaction time τ_{BA} . Its error was therefore of little impact. On the other side, the collisions $N_{\text{coll}}(r)$ with the central target, which form the main contribution to Eq. (21), were much less affected by the chain distribution far outside the center (Figure 2, right) and were extrapolated accurately, despite of the sparse chain density at the cell periphery. This feature contributes to the fact that the extrapolation process appears to be insensitive to the chain distribution far away from the target. Similarly, Eq. (22) delivered consistent and accurate results, except for the last data point which belongs




FIG. 4: Extrapolation of the reaction time τ_{BA} to large cell radii R_L . The dotted curve is the analytical estimate Eq. (11), the solid and dashed curves correspond to the MEC-E equations (21) and (22), respectively. Upper triple: $p = 1$ (free diffusion). Lower triple: $p = 2^{-8}$, where facilitated diffusion is most effective.

to the highest walker-chain affinity. Here, the sliding-length already reaches one half of the system size that was used to fit the empirical parameters. A larger dimensioned test system is required for such high affinities to increase the accuracy of the extrapolation procedure.

The simulation time required to set up the MEC-E equations (21) and (22) equals the average number of time steps the walker needed to reach the radius $R_S = 4$ when starting at the central target, i.e. $\tau_f(p, R_S)$. Compared to the corresponding time required to simulate $\tau_{BA}(R_L)$ explicitly, a speed up between 50 and 500 was gained, depending upon walker-chain affinity (Table I, last column). Integrated over all 12 data points, a total speed up of 140 was derived.

It is possible and intended to exploit this method for extrapolations to much larger systems. Figure 4 displays the extrapolation of $\tau_{BA}(p, R_L)$ up to $R_L = 20$ for $p = 1$ (free diffusion) and $p = 2^{-8}$, close to the minimum in Figure 3. The chain density was assumed to remain constant, i.e. its length was growing as $L(R_L) \sim R_L^3$. Explicit simulations of τ_{BA} are not feasible any more for such large cells. However, for free diffusion, Eq. (10) is available, and both extrapolation methods delivered reaction times about 8% above the exact solution, which, in this plot, was un-distinguishable from the approximation Eq. (15). When protein-chain interaction was enabled, both extrapolation methods delivered almost identical results, which were about 15% above the analytical estimate Eq. (11).

7. SUMMARY

In this work, the empirical MEC-E equations (21) and (22) were derived and tested against random walk simulations. Whereas the original MEC approach (Sect. 2.2) represents an exact method to obtain the average reaction time τ_{BA} by simulating the much faster back-reaction $\mathbf{A} \rightarrow \mathbf{B}$, it still requires to set up a model system of full size R . This would become prohibitive in simulations of large cells containing realistic chains with thermal fluctuations and hydrodynamic interactions.

We have demonstrated that the simulation of a test system of moderate size is sufficient to extract reaction times of much larger cells. This is so because the number of collisions as a function of the cell radius, $N_{\text{coll}}(r)$, is asymptotically approaching a plateau (Figure 2, right). In this region, the reaction time is merely proportional to the effective volume V_{eff} , as shown in Eq. (21), with a small correction in form of the first passage time $\tau_f(r)$, Eq. (17). This quantity is easily estimated once the effective diffusion coefficient is determined. If the test system is too small for $N_{\text{coll}}(r)$ to reach the plateau, it is still possible to obtain accurate results, because the functional form of this quantity is known (Eq. 18 and 19), so that extrapolations to larger cells become feasible.

The size of the test system has to be chosen with care, because only those regions are of use in which the walker experiences a randomized and approximately homogeneous environment. Within the central region, typically of the size of the sliding length ξ , the reaction time is dominated by 1-dimensional (sliding) instead of 3-dimensional diffusion. This part of the cell has to be excluded when the walker's diffusion properties are analyzed. The same holds true for the outermost region, where the chain conformation exhibits boundary effects. Assuming that the sliding length ξ does not exceed the persistence length l_p , a cell radius R of five persistence lengths appears adequate. Here, the region $\xi < r < R - 2l_p$ may be exploited to set up the empirical equations (21) or (22). With increasing walker-chain affinity and sliding length, the radius R has to be adjusted accordingly.

The results presented above demonstrate how the MEC-E approach delivers a speed up between 50 and 500 (depending on walker-chain affinity, Table I) by extrapolation from $R_S = 4$ to $R_L = 6$, with respect to explicit simulations of the reaction time τ_{BA} . With increasing radius R_L , Eq. (22) is approximated as

$$\tau_{\text{Sz,eff}}(R_L \gg R_{\text{eff}}) \approx \frac{R_L^3}{3 D_{\text{eff}} R_{\text{eff}}}, \quad (23)$$

and the speed up is therefore approximately growing proportional to R_L^3 .

APPENDIX A: PROOF OF EQUATION (18)

As was shown by Berg [14], the probability of a walker, after starting at r_{ini} (where $R_a < r_{\text{ini}} < R$), to be adsorbed at R_a , before reaching the distance R , is

$$P(R) = \frac{R_a(R - r_{\text{ini}})}{r_{\text{ini}}(R - R_a)}. \quad (A1)$$

This was derived from the steady-state solution of Fick's second equation for spherical symmetry,

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{dC(r)}{dr} \right) = 0. \quad (A2)$$

Here, $C(r)$ is the concentration, having a maximum at the particle source radius $r = r_{\text{ini}}$ and dropping to zero at the adsorbers radii $r = R_a$ and $r = R$.

In our case, not the probability $P(r)$, but the average number $N_{\text{coll}}(r)$ of events in which the walker returns to $r = R_a$ before first reaching the distance $r = R$ is of interest. We shall now assume that $N_{\text{coll}}(r)$ is known for one particular distance r , and we want to derive $N_{\text{coll}}(r + dr)$. The probability, that the walker, starting from r , goes straight to $r + dr$, is $1 - dP(r)$. Then, the probability to first return back to the target, before passing through r and reaching $r + dr$, is $dP(1 - dP)$. In this latter case, $2 \cdot N_{\text{coll}}(r) + 1$ collisions have already occurred in average. The probability to return exactly n times to the target and back to r before reaching $r + dr$ is $dP^n(1 - dP)$, yielding $(n + 1) \cdot N_{\text{coll}}(r) + n$ collisions. The sum

$$\begin{aligned} N_{\text{coll}}(r + dr) &= (1 - dP) \sum_{n=1}^{\infty} [n N_{\text{coll}}(r) + n - 1] dP^{n-1} \\ &= \frac{N_{\text{coll}}(r) + 1}{1 - dP} - 1 \end{aligned} \quad (A3)$$

leads to the differential equation

$$\frac{dN(r)}{dr} = N(r) \frac{dP}{dr} + \frac{dP}{dr}. \quad (A4)$$

With Eq. (A1) we further have

$$dP = \frac{R_a dr}{r(r - R_a)}, \quad (A5)$$

so that Eq. (A4) is solved as

$$N_{\text{coll}}(r) = (N_{\infty} + 1) \frac{(r - R_a)}{r} - 1. \quad (A6)$$

Here, $N_{\infty} = N_{\text{coll}}(r \rightarrow \infty)$ is the asymptotic limit for the number of collisions far away from the target. This solution is incorrect close to the target, where $N_{\text{coll}}(R_a) = -1$. In fact, the validity of this approach is restricted to length scales that are large compared to the (finite) step-size. In particular, since we want extrapolate $N_{\text{coll}}(r)$ to

a large distance, we can assume r to be large enough so that $N_{\text{coll}}(r) \gg 1$. Then, the sum Eq. (A3) simplifies to

$$\begin{aligned} N_{\text{coll}}(r + dr) &= (1 - dP) \sum_{n=1}^{\infty} n N_{\text{coll}}(r) dP^{n-1} \\ &= \frac{N_{\text{coll}}(r)}{1 - dP}, \end{aligned} \quad (\text{A7})$$

leading to

$$\frac{dN(r)}{dr} = N(r) \frac{dP}{dr}, \quad (\text{A8})$$

which finally solves to

$$N_{\text{coll}}(r) = N_{\infty} \frac{(r - R_a)}{r}. \quad (\text{A9})$$

Both parameters N_{∞} and R_a were used as free fit parameters. We have verified that Eq. (A6) and Eq. (A9) deliver identical results when extrapolating to large radii, so that, for sake of simplicity, Eq. (A9) was applied throughout this work.

APPENDIX B: PROOF OF EQUATION (19)

When considering Eq. (1),

$$\tau_{\text{BA}} + \tau_{\text{AB}} = (N_{\text{coll}} + 1) \cdot \tau_R,$$

we note that in case of free diffusion the reaction time τ_{BA} is given by Eq. (10) and τ_{AB} by Eq. (17) with the free diffusion coefficient D , Eq. (13), so that

$$(N_{\text{coll}}(r) + 1) \cdot \tau_R(r) = f(r) \quad (\text{B1})$$

and

$$f(r) = \frac{r^2}{3D} \cdot \left(\frac{r}{R_a} + \frac{R_a^2}{2r^2} - 1 \right) \quad (\text{B2})$$

with $r > R_a$. Using Eq. (2) we obtain

$$N_{\text{coll}}(r) = \frac{f(r)}{\tilde{\tau}_R V_{\text{eff}}(r)} - 1. \quad (\text{B3})$$

Both quantities D and the effective source radius R_a are used as free fit parameters.

-
- [1] A.D. Riggs, S. Bourgeois and M. Cohn, *The lac repressor-operator interaction. 3. Kinetic studies*, J. Mol. Biol. **53**, 401 (1970).
 - [2] P.H. Richter and M. Eigen, *Diffusion controlled reaction rates in spheroidal geometry. Application to repressor-operator association and membrane bound enzymes*, Biophys. Chem., **2**, 255 (1974).
 - [3] O.G. Berg and P.H. von Hippel, *Diffusion-controlled macromolecular reactions*, Annu. Rev. Biophys. Chem. **14**, 130 (1985).
 - [4] M. Ptashne and A. Gann, *Genes and Signals*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. (2001).
 - [5] O.G. Berg, R.B. Winter and P.H. von Hippel, *Diffusion driven mechanisms of protein translocation on nucleic acids. 1. Models and theory*, Biochemistry **20**, 6929 (1981).
 - [6] H.X. Zhou and A. Szabo, *Enhancement of Association Rates by Nonspecific Binding to DNA and Cell Membranes*, Phys. Rev. Lett. **93**, 178101 (2004).
 - [7] S.E. Halford and J.F. Marko, *How do site-specific DNA-binding proteins find their targets?*, Nucleic Acids Research **32**, 3040 (2004).
 - [8] K. Klenin, H. Merlitz, J. Langowski and C.X. Wu, *Facilitated diffusion of DNA-binding proteins*, Phys. Rev. Lett. **96**, 018104 (2006).
 - [9] Tao Hu, A.Yu. Grosberg, B.I. Shklovskii, *How do proteins search for their specific sites on coiled or globular DNA*, arXiv:q-bio.BM/0510043 (2005).
 - [10] H. Merlitz, K. Klenin, C.X. Wu and J. Langowski, *Facilitated diffusion of DNA-binding proteins: Efficient simulation with the method of excess collisions (MEC)*, J. Chem. Phys. **124** (2006) (in print).
 - [11] K.V. Klenin and J. Langowski, *Modeling of intramolecular reactions of polymers: An efficient method based on Brownian dynamics simulations*, J. Chem. Phys. **121**, 4951 (2004).
 - [12] A. Szabo, K. Schulten and Z. Schulten, *First passage time approach to diffusion controlled reactions*, J. Chem. Phys. **72**, 4350 (1980).
 - [13] K. Klenin, H. Merlitz and J. Langowski, *A Brownian Dynamics Program for the Simulation of Linear and Circular DNA and other Wormlike Chain Polyelectrolytes*, Biophys. J. **74**, 780 (1998).
 - [14] Howard C. Berg, *Random walks in Biology*, Princeton University Press, expanded edition (1993).